

Chapter 4

No Peeking

Why is interference destroyed by “which path” information? And if we only obtain partial information, is the interference only degraded not completely destroyed?

Last Update: 30/12/11

1. Two Slit Experiment and Perfect “Which Path” Information

Consider the familiar two-slit arrangement. An interference pattern is developed on the screen. In terms of the Hilbert space states this can be understood as follows. The state of the wave/particle prior to hitting the screen is a coherent superposition of the two states representing passage through each slit, thus,

$$|\psi\rangle = \frac{1}{\sqrt{2}}(|U\rangle + |L\rangle) \quad (1)$$

where U and L stand for the upper and lower slits. Each of the states $|U\rangle$ and $|L\rangle$ is normalised to unity, representing a single particle. Consider a given position, x , on the screen. Coherence means that we can consider the upper wave differing from the lower only by some phase difference which depends upon x , say $\theta(x)$, so that

$|U\rangle = e^{i\theta(x)}|L\rangle$. Consequently (1) becomes,

$$|\psi\rangle = \frac{1}{\sqrt{2}}(e^{i\theta(x)} + 1)|L\rangle \quad (2)$$

The intensity of the pattern at position x on the screen is proportional to $|\psi|^2 = \langle\psi|\psi\rangle$ which (2) gives to be,

$$\langle\psi|\psi\rangle = 1 + \cos(\theta(x)) \quad (3)$$

Equ.(3) produces the familiar interference pattern, with alternating dark bands when θ is an odd multiple of π (zero intensity), and bright bands where θ is an even multiple of π (intensity 2 units). The average intensity is unity, as it should be to consistently represent a single particle.

Now let us introduce a two-state device, M , in the beam path of the lower slit. This device is set up in state $|M : U\rangle$ but it will change to state $|M : L\rangle$ if a particle passes through it. In other words M is a device which records which path the particle takes.

Readers will recall that such a measurement is expected to destroy the interference pattern. But why should this always be the case – for any sort of measurement, M , however we choose to contrive it? The impression that is sometimes given is that the wave/particle is physically disturbed by the measurement, and that it is this physical disturbance which destroys the interference. This is seriously misleading and misrepresents the nature of quantum mechanics. The vanishing of the interference arises from the very possibility of distinguishing paths and is a result of the state algebra alone (or, if you will, the quantum logic). The means of obtaining the “which path” information, however subtle, is irrelevant. This will now be demonstrated.

Suppose that M is a perfect measuring device, and suppose that we are certain that exactly one particle has passed through our apparatus. This means that M will

infallibly switch to state $|M : L\rangle$ if the particle takes the lower path and infallibly be left in state $|M : U\rangle$ if the particle takes the upper path. However, to be a perfect measurement we must be able to distinguish with certainty between the two states $|M : L\rangle$ and $|M : U\rangle$. In other words, these two states must be orthogonal,

$$\text{Perfect Measurement:} \quad \langle M : L | M : U \rangle = 0 \quad (4)$$

Prior to any interaction between our particle and the measuring device, their combined state is just the direct product state $|\Psi\rangle = |\psi\rangle|M : U\rangle = \frac{1}{\sqrt{2}}(|U\rangle + |L\rangle)|M : U\rangle$ because M is set up in state $|M : U\rangle$. However, after the wave/particle has had opportunity to interact with M the state evolves into,

$$|\Psi\rangle \rightarrow |\Psi'\rangle = \frac{1}{\sqrt{2}}(|U\rangle|M : U\rangle + |L\rangle|M : L\rangle) \quad (5)$$

The intensity of the signal received on the screen is proportional to $\langle\Psi'|\Psi'\rangle$ which, from (5), is now simply,

$$\langle\Psi'|\Psi'\rangle = 1 \quad (6)$$

i.e., uniform illumination with no interference fringes. Why? Because the orthogonality of $|M : L\rangle$ and $|M : U\rangle$ causes the two terms on the RHS of (5) to be orthogonal. The cross-product between them is now zero, whereas previously it was the cross-product which gave rise to the $\cos\theta$ term in (3) and hence the interference.

It is clear, then, that the nature of the measurement, M , is irrelevant. The interference disappears by virtue of the perfect nature of the measurement as expressed by the orthogonality of the measurement state, (4).

2. Imperfect “Which Path” Information

We can now see that the destruction of the interference pattern need not be an all-or-nothing affair. Suppose that M is an imperfect measuring device which has $\langle M : L | M : U \rangle \neq 0$. Then,

$$\langle\Psi'|\Psi'\rangle = 1 + \Re\left(e^{i\theta(x)}\langle M : L | M : U \rangle\right) \quad (7)$$

Consequently the interference pattern is merely dimmed if $0 < |\langle M : L | M : U \rangle| < 1$ but does not vanish completely.

In the early days of quantum mechanics, great emphasis was placed on measurements requiring physical interaction with the measured system and the (supposed) impossibility of making measurements without disturbing what was being measured. Such physical disturbance was held to be responsible for the destruction of the interference in cases like that considered above. However, this is demonstrably incorrect. A graphic illustration of this is provided by the Elitzur-Vaidman bomb test. This involves a bomb whose sensitivity is so extreme that it is guaranteed to explode if exposed any physical interaction whatsoever, however minor. If this bomb is used as a measuring device it can successfully eliminate an interference pattern without exploding. (This is discussed in detail in [Chapter 8](#)). So elimination of the interference is clearly not caused by physical disturbance since any actual physical disturbance

would have resulted in the bomb exploding. What *is* necessary is the *possibility* of physical interaction.

Another way of appreciating this point is to note that the evolution of the state into the form of Equ.(5) does not really constitute a measurement. It is merely the first part of a measurement process. The second part is the ‘collapse of the wavefunction’ in which only one or other of the two possibilities, either $|U\rangle|M : U\rangle$ or $|L\rangle|M : L\rangle$, is actually realised. But if the combined particle-plus- M system is unperturbed by interaction with any third party, including our own selves, then their combined state remains as given in (5). But this state can be obtained by a purely unitary evolution. In other words, $|\Psi\rangle \rightarrow |\Psi'\rangle$ is calculable from the Schrodinger equation, so that

$$|\Psi'\rangle = \exp\left\{-\frac{i}{\hbar}\hat{H}t\right\}|\Psi\rangle \text{ where } \hat{H} \text{ is the appropriate Hamiltonian operator.}$$

Consequently it is possible, by another unitary evolution, for (5) to be converted back again, clearly so because $|\Psi\rangle = \exp\left\{\frac{i}{\hbar}\hat{H}t\right\}|\Psi'\rangle$. This is known as “quantum erasure”

and will be considered further in [Chapter 18](#). The quantum erasure process re-establishes an interference pattern which would otherwise have been destroyed by the interaction with M . This would not be possible for a genuine measurement since the collapse of the wavepacket is irreversible.

3. Three Slits and Partial Information

We have seen that an imperfect measurement on two slits causes degrading of the interference rather than complete elimination. What happens if we use three slits but obtain information from only one?

In obvious notation, and with no measuring device, the state is,

$$|\psi\rangle = \frac{1}{\sqrt{3}}(|1\rangle + |2\rangle + |3\rangle) \quad (8)$$

Suppose the relative phase between the waves from slits j and k is $\theta_{jk}(x)$ at position x on the screen, then there is an interference pattern on the screen given by,

$$\langle\psi|\psi\rangle = 1 + \frac{2}{3}[\cos\theta_{12}(x) + \cos\theta_{23}(x) + \cos\theta_{13}(x)] \quad (9)$$

For example, if $\theta_{12} = \theta_{23} = \theta$ and $\theta_{13} = 2\theta$ then,

$$\langle\psi|\psi\rangle = 1 + \frac{2}{3}[2\cos\theta(x) + \cos 2\theta(x)] \quad (10)$$

The average is unity, as it must be, and the intensity varies from 0 to 3.

Now, suppose we introduce our two-state device M into the path of slit 3 (only). This does not provide a measurement which discriminates between slits 1 and 2. So does a modified interference pattern remain, namely that due to slits 1 and 2 alone? The answer is “yes” as may be seen by considering the state algebra as follows. The combined state evolves thus,

$$|\Psi\rangle \rightarrow |\Psi'\rangle = \frac{1}{\sqrt{3}}(|1\rangle|M : U\rangle + |2\rangle|M : U\rangle + |3\rangle|M : L\rangle) \quad (11)$$

Here $|M : U\rangle$ is the state in which M is initially setup, and hence it remains in this state if the particle takes paths 1 or 2. Only path 3 results in the modified M state, $|M : L\rangle$. Consequently when we form the absolute square of the combined state, which provides the intensity seen at the screen, the third term in (11) is orthogonal to the first two, but the first two terms continue to provide an interference cross-product,

$$\langle \Psi' | \Psi' \rangle = 1 + \frac{2}{3} \cos \theta(x) \quad (12)$$

Hence (12) continues to show an interference pattern, though with diminished contrast between the darker and lighter bands compared with that obtained with no measuring device. The minima are now $1/3$ and the maxima $5/3$, compared with 0 and 3 when all three slits contribute. The average is again unity, as it must always be. This is a further illustration of how partial information will degrade the interference pattern without removing it completely.

This document was created with Win2PDF available at <http://www.win2pdf.com>.
The unregistered version of Win2PDF is for evaluation or non-commercial use only.
This page will not be added after purchasing Win2PDF.